



Volume 27, Issue 10, Page 587-595, 2024; Article no.JABB.123643 ISSN: 2394-1081

# Scanning Electron Microscopy Based Morphology of Papaya Mealybug [*Paracoccus marginatus* Williams and Granara De Wilink (Hemiptera: Pseudococcidae)] on Mulberry Plant of Assam, India

# M. Hazarika <sup>a\*</sup>, S. K. Dutta <sup>b</sup>, B. Bhattacharryya <sup>b</sup> and D. Dey <sup>c</sup>

<sup>a</sup> College of Sericulture, Assam Agricultural University, Jorhat-78013, Assam, India.
<sup>b</sup> Department of Entomology, Assam Agricultural University, Jorhat-78013, Assam, India.
<sup>c</sup> Division of Entomology, Indian Agricultural Research Institute, New Delhi, India.

#### Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

#### Article Information

DOI: https://doi.org/10.9734/jabb/2024/v27i101481

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: https://www.sdiarticle5.com/review-history/123643

> Received: 22/07/2024 Accepted: 24/09/2024 Published: 30/09/2024

Original Research Article

#### ABSTRACT

A scanning electron microscopy (SEM) study was conducted on the external morphology of the adult female Papaya mealybug, *Paracoccous marginatus* Willam and Granara de Willink. This study aimed to enhance the understanding of the external morphology and wax-secreting pores of the papaya mealybug through SEM examination. The ultrastructure of the antenna, stylet fascicle,

\*Corresponding author: E-mail: manaseehazarika104@gmail.com, manasee.hazarika@aau.ac.in;

*Cite as:* Hazarika, M., S. K. Dutta, B. Bhattacharryya, and D. Dey. 2024. "Scanning Electron Microscopy Based Morphology of Papaya Mealybug [Paracoccus Marginatus Williams and Granara De Wilink (Hemiptera: Pseudococcidae)] on Mulberry Plant of Assam, India". Journal of Advances in Biology & Biotechnology 27 (10):587-95. https://doi.org/10.9734/jabb/2024/v27i101481.

legs, cerarii, vulva, and wax-secreting pores was carefully examined. Despite the time-consuming and expensive nature of preparing specimens for SEM, the detailed information obtained is valuable for resolving challenging aspects of papaya mealybug taxonomy and enhancing our understanding of the role of wax in their biology.

Keywords: SEM; Papaya mealybug; adult female; morphological characters.

# 1. INTRODUCTION

"The Papaya mealybug (PMB) Paracoccous marginatus Willam and Granara de Willink (Hemiptera: Pseudococcidae) is a troublesome insect pest known to infest various economically important plants such as papaya, mulberry, acalypha, hibiscus, plumeria, avocado, citrus, cotton, tomato, eggplant, pepper, beans and peas, sweet potato, mango, cherry and pomegranate" [1,2]. This insect caused heavy infestation and yield reduction of mulberry throughout India recently. "It is an exotic polyphagous-sucking insect with high multiplication and spreading potential at the field level. It is native to Mexico and Central America. where it never acquires the status of a serious pest, probably due to the presence of an endemic natural enemy complex" [3]. Though its presence was known since 1955, "it was only described in 1992" [1]. "PMB became a pest when it invaded the Caribbean region. Since 1994 it has been recorded in 14 Caribbean countries. The pest was recorded in Bradenton, Florida in 1998 on Hibiscus, and by 2002 it spread to 18 different plant species in 30 different cities. The establishment of this pest in Guam in 2002" [4] and "Palau in 2003 resulted in further spread to neighboring Hawaiian Islands in the Pacific. In Asia, it was reported from Indonesia, India, and Sri Lanka in 2008" [5], Maldives and Bangladesh in 2009 and Thailand, Cambodia and the Philippines in 2010. "It was also reported from Reunion Island in the Arabian Sea Ghana in West Africa in 2010 and Assam in 2013" [6].

The SEM is a valuable tool for studying the detailed morphology of small structures on insect cuticles, and in the case of mealybugs, it helps in understanding the significance of waxes produced by different dermal pores in protecting developmental stages. The rigid exoskeleton of insects is especially suitable for electron microscopy because the specimen usually retains its shape under vacuum. "Wax on the body surface may prevent contamination by honeydew" [7,8,9] or "invasion of the body by bacteria, fungi, and other parasites" [10] or

"prevent damage at the point where eggs touch, so facilitating hatching" [11]. "The mealy wax covering on the integument of mealybugs differs from that of other insects in many ways, its presence on the surface is well documented but there has been only limited speculation on its function or reasons for the variety of pores and ducts found within a single species" [12,13]. "The structure of the wax-producing pores is complicated and can be elucidated better using SEM. However, only a few studies have reported ultrastructure and wax-secretion on the processes in the family Pseudococcidae" [8,14]. Likewise, a few studies reported on the ultrastructure and wax-secretion processes in the family Pseudococcidae. There has not been any study of the structure of Papaya mealybug, hence, this study aimed to fill the gap in the understanding of the external morphology and wax-secreting glands of the Papaya mealybug through scanning electron microscopy.

# 2. MATERIALS AND METHODS

The scanning electron microscopic studies focused on the adult female, as the external micrographic characters of this stage are considered taxonomically vital. Adult females were kept in 100% acetone for 1 hour. The overnight 0.25% sample was kept in Glutaraldehyde fixative in the refrigerator at 4°C. After overnight keeping in fixative, samples were then washed three times in 0.1M phosphate buffer and were dehydrated by passing through a series of 30, 50, 70, 90, and 100% alcohol for 15 min in each concentration. After that 1 drop of Hexamethyldisilazane (HMD) with 1 drop of alcohol on the sample was open-dried. Each dehydrated specimen was then glued onto a standard SEM stub (diameter 12.5 mm, height 3.0 mm) using double-sided sticky tape. SEM stubs were then put in a sputter clouter for 90 minutes. To prevent the electron beam from burning a hole through the cuticle, each specimen was coated with a protective, reflective surface. Sputter coating covers the specimen with a very thin layer of heavy metal (goldpalladium mixture). Each specimen was placed in the sputter chamber, the air was pumped out and replaced with nitrogen gas, before which a plasma of metal molecules was released for 3.5-4.0 minutes to coat the specimen with metal. A fully computer-controlled scanning electron microscope ZEISS (EVO MA10) with a tungsten heated filament was used to scan each gold-palladium-coated specimen.

#### 3. RESULTS

The scanning electron micrographic studies revealed detailed external morphological characters of the adult female Papaya mealybug *viz.*, antenna, stylet fascicle, legs, cerarii, vulva, and wax-secreting pores revealed the following-

Antenna: Eight antennal segments were present in adult female papaya mealybug. The apical (flagellum) and sub-apical (scape and pedicel) segments clearly separated. Scape was a strong and stout segment followed by a less stout pedicel. Pedicel had 4 recurvate setae (3 on the dorsomedial aspect and 1 on the ventromedial aspect). The total number of the flagellar segment was eight, the apical most flagellar segment is the longest and the ventrolateral aspect of the flagellar segments had 10–12 flagellate setae, 3 fleshy setae, and 6 stiff setae near the apex. Two comparatively stouter and more pointed setae were present on the ventrolateral aspect (Fig. 1). All the antennal setae appeared to be trichoid sensillae for olfactory, gustatory, and thigmotactic functions.

**Stylet fascicle:** Stylet fascicle consisted of 4 distinct stylets (2 mandibular and 2 maxillary), labrum and labium were fused to form the proboscis. Setae on the proboscis was eract with curved tips These setae appeared to be chemosensillae for olfactory and gustatory response (Fig. 2).

**Legs:** Adult female mealybugs had only one tarsal segment and a single claw on the tarsus of each leg. At the expanded base of the claw, there was usually a pair of seta-like digitules (claw digitules), which were either shorter or longer than the claw, and either pointed or expanded distally. Coxa was the most robust segment with 2-3 recurvate setae, the trochanter is triangular, ventromedial seta in the trochanter was the longest. The femur was the longest segment with 4 setae in ventro-medial and 3 setae in dorso-medial aspects. In the tibia, 3 stout setae were present in the ventromedial aspect. Tarsus uni-segmented as described (Fig. 3).





Hazarika et al.; J. Adv. Biol. Biotechnol., vol. 27, no. 10, pp. 587-595, 2024; Article no.JABB.123643



Fig. 2. Stylet fascicle of female P. marginatus

**Cerarii:** Each cerarius usually contains more than two conical setae and an aggregation of trilocular pores (Fig. 7).

**Pygidium:** Pygidium was heavy with an inwardly rounded periphery. Long and fleshy setae were present in pygidium apex of pygidium with, 4-5 inwardly curved long setae. Discoidal pores are present on the dorsal surface of the pygidial body (Fig. 4).

**Vulva:** The elevated, triangular vulva has a "T"-shaped slit opening with striations that radiate outward (Fig. 5).

**Wax-secreting pores:** The wax-secreting pores of adult female *P. marginatus* were as follows:

a. Multilocular pores: They are occasionally seen on the dorsum, although they are typically lacking on the venter, at least around the vulva. All of the pores were made up of an outer ring with 8–10 loculi and a circle with more than five holes. Each egg developed an

attachment to the wax curls created by the multilocular pores surrounding the vulva."The wax curls produced by the multilocular pores surrounding the vulva became attached to each egg as it left the genital opening" [14]. Wax produced by multilocular disc pores envelops the eggs in the ovisac, shielding them from natural enemies. precipitation, desiccation. and honeydew contamination. After their second instar, males also employ wax from the multilocular disc pores to build their cocoons (Fig. 6).

b. Trilocular pores: "Trilocular pores produce spiral wax filaments that fragment to form the fine, mealy body covering [8]. These pores were usually abundant and evenly distributed over both dorsum and venter; each consisted of a rounded-triangular, raised area having three elongate, narrow openings radiating from the center of the pore and slightly twisted, or swirled, about each other" [8] (Fig. 7). Hazarika et al.; J. Adv. Biol. Biotechnol., vol. 27, no. 10, pp. 587-595, 2024; Article no.JABB.123643



c. Hindleg

Fig. 3. a. Foreleg, b. middleleg, c. hindleg of female P. marginatus



Fig. 4. Pygidium of female P. marginatus



Fig. 5. 'T' shaped slit of vulva of female P. marginatus



Fig. 6. Multilocular pore of female P. marginatus



Fig. 7. Trilocular pore of female P. marginatus

#### 4. DISCUSSION

The results of the study contribute to the understanding of the external morphology of the Papaya mealybug, providing valuable information for taxonomic identification and differentiation from related species. "From the experiment, micrographs of the papaya mealybug's external morphological characters (antenna, stylet fascicle, legs, cerarii, vulva, and pores that secrete wax) were not made by anyone else, but comparable research was conducted on six specific mealybug species found in Sri Lanka" (Sirisena et. al., 2015). "Early workers used the number of antennal setae as primary characters to separate species" [15,16,17]. "The overall length of the antenna, as well as the length of the individual segments, were recorded in detail by numerous workers until Ferris [18] decided that the variation made was unreliable as a taxonomic character". "Differences in wax pore ultrastructure may be taxonomically useful. Such differences have been used to revise the generic placement of some scale insect species into families, such as Coccidae, Diaspididae, and Monophlebidae" [19,20,21,22]. "Adult female mealybugs are covered with hundreds of pores most of which produce various types of wax for different purposes" [23,24]. "These pores are distinct sclerotized structures that act as molds to produce structurally different forms of wax in different areas of the body" [24]. "The wax curls

produced by the multilocular pores surrounding the vulva became attached to each egg as it left the genital opening" [14]. The details of the setae and pores recorded in the present study would be useful in ascertaining the identity of the species and differentiating it from other related species. Moreover, it will provide a basis for the secretion of the pores and their significance in the biology of the pest.

#### **5. CONCLUSION**

The scanning electron microscope (SEM) is a crucial tool for studying the detailed morphology of very small structures on insect cuticles, providing quick and accurate representation. SEM is a very useful tool for observing minute, three-dimensional structures of papaya mealybug, and some of these details can only seen under the SEM. Despite the timeconsuming and expensive nature of preparing specimens for SEM, the detailed information obtained is essential for resolving challenging aspects of papaya mealybug taxonomy and improving understanding of the role of wax in their biology.

#### **DISCLAIMER (ARTIFICIAL INTELLIGENCE)**

Author(s) hereby declare that NO generative AI technologies such as Large Language Models (ChatGPT, COPILOT, etc.) and text-to-image

generators have been used during the writing or editing of this manuscript.

#### ACKNOWLEDGEMENT

The author expresses gratitude to Head, Department of Entomology, IARI, New Delhi for having kindly given the opportunity to carry out the Scanning Electron Microscopic work in the departmental laboratory. The authors express gratitude and indebtedness to Department of Science and Technology, Government of India for financial assistance in terms of DST-INSPIRE Fellowship to the author.

### **COMPETING INTERESTS**

Authors have declared that no competing interests exist.

# REFERENCES

- Williams DJ, Granara de Willink MC. Mealybugs of Central and South America. CAB International, Wallingford, England. 1992;635.
- 2. Miller DR, Miller GL. Redescription of Paracoccus marginatus Williams and De Willink (Hemiptera: Granara Coccoidea: Pseudococcidae) including descriptions of the immature stages and adult male. Proc. Entomol. Soc 2002:104:123.
- 3. Tanwar RK, Jeyakumar P, Vennila S. Papaya mealybug and its management strategies. Tech. Bull. 22. NCIPM, New Delhi. 2010;22.
- Meyerdirk DE, Muniappan R, Warkentin R, Bamb J, Reddy GVP. Biological control of the papaya mealybug, *Paracoccus marginatus* (Hemiptera: Pseudococcidae) in Guam. Plant Protect. Qtly. 2004;19(3).
- Muniappan R, Shepard BM, Watson GW, Carner GR, Sartiami D, Rauf A, Hammig MD. First report of the papaya mealybug, *Paracoccus marginatus* (Hemiptera: Pseudococcidae), in Indonesia and India. J. Agric. Urban Entomol. 2009;25(1):37-40.
- 6. Sarma AK. Invasion of papaya mealybug, *Paracoccus marginatus* in Assam. Indian J. Entomol. 2013;75:355-356.
- 7. Broandbent L. Aphid excretion. Poc. R. Entomol. Soc. Lond. 1951;26:97-103.
- Cox JM, Pearce MJ. An experimental study of morphological variation in mealybugs (Homoptera: Coccidae: Pseudococcidae). Syst. Entomol.1983;8:361-382.

- 9. Pope RD. Some aphid waxes, their form and function (Homoptera: Aphididae). J. Nat. Hist. 1983;17:489-506.
- Foldi I, Pearce MJ. Fine structure of wax glands, wax morphology and function in the female scale insect, *Pulvinaria regalis* Canard (Hemiptera: Coccidae). Int. J. Insect Morphol. Embryol. 1985;14:259-271.
- 11. Hashimoto A, Ueda M. Coil form wax on the egg surface of *Drosicha corpulenta* Kuwana (Homoptera: Margarodidae). Appl. Entomol. Zool. 1985;20:92-93.
- 12. Waku Y, Manabe Y. Fine structure of the wax gland in a scale insect, *Eriococcus lagerstraemiae* Kuwana (Homoptera: Eriococcidae). Appl. Entomol. Zool. 1981;16(02):94-102.
- Foldi I, Cassier P. Ultrastructure comparée des glands tegumentaires de treize families de cochenilles (Homoptera: Coccoidea). Int. J. Insect Morphol. Embryol. 1985;14:33-50.
- Kumar V, Tewari SK, Datta RK. Dermal pores and wax secretion in mealybug *Maconellicoccus hirsutus* (Hemiptera, Pseudococcidae). A pest of mulberry. Ital. J. Zool. 1997;64(4):307-311.
- Maskell WM. Further coccid notes with descriptions of several new species and discussion of various points of interest. Trans. Proc. New Zeal. Inst. 1894;26:65-105.
- 16. Essig EO. The genus *Pseudococcus* in California. Pomona Coll. J. Entomol. 1909;1:35-46.
- Hollinger AH. Taxonomic value of antennal segments of certain Coccidae. Ann. Entomol. Soc. America. 1917;10: 264-271.
- Ferris GF. The alleged occurrence of a seasonal dimorphism in the females of certain species of mealy bugs (Hemiptera; Coccidae). Entomol., News. 1918;29:349-352.
- 19. Takagi S. Disc pores of Diaspididae: Microstructure and taxonomic value (Homoptera: Coccoidea). Insecta Matsumurana. 1990;44:81-122.
- Ülgentürk S, Wilhem G. Wax exuding discpores of *Coccus hesperidum* L. *Parthenolecanium corni* Bouché and *Sphaerolecanium prunastri* (Fonscolombe) (Hemiptera: Coccidae), In: Pellizzari, G. (Ed.). Programme and Abstracts. IX International Symposium on Scale Insect Studies, Padua, Italy. 2001;82.

- Unruh CM. A taxonomic review of the *Crypticerya* species (Hemiptera: Coccoidea: Monophlebidae) of the southwestern United States and Mexico, including description of a new species from Baja California. Zootaxa. 2008;1759: 1-42.
- 22. Unruh CM, Gullan PJ. Molecular data reveal convergent reproductive strategies in iceryine scale insects (Hemiptera: Coccoidea: Monophlebidae), allowing

reinterpretation of morphology and a revised generic classification. Syst. Entomol. 2008;33:8-50.

- 23. McKenzie HL. Mealybugs of California. University of California Press, LosAngeles. 1967;526.
- 24. Cox JM, Pearce MJ. Wax produced by dermal pores in three species of mealybug (Homoptera: Pseudococcidae). Int. J. InsectMorphol. Embryol. 1985;12: 235-248.

**Disclaimer/Publisher's Note:** The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/123643