



The Nutraceuticals: A Voluminous Torrent in Pharmaceuticals-coupling Health & Drugs

Vijay Sharma^{1*}, Lalit Singh¹ and Navneet Verma²

¹Department of Pharmaceutics, SRMSCET, Bareilly, U.P., India.

²Faculty of Pharmacy, IFTM University, Moradabad, U.P., India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMPS/2016/25467

Editor(s):

(1) Dongdong Wang, Department of Pharmacognosy, West China College of Pharmacy, Sichuan University, China.

Reviewers:

(1) Francesco Marotta, Texas Women University, USA.

(2) Jose Juis Montañez Soto, Instituto Politécnico Nacional, Mexico.

(3) Josephine Ezekwesili-Ofili, Nnamdi Azikiwe University, Nigeria.

Complete Peer review History: <http://sciencedomain.org/review-history/14624>

Review Article

Received 4th March 2016
Accepted 22nd April 2016
Published 14th May 2016

ABSTRACT

The present review is focused on the nutraceuticals which are present in our surroundings having an excellent impact over the health of humans but are not known for their pharmaceutical use. Nutraceuticals include vitamins, minerals and other dietary supplements, which may be herbs, enzymes, animal extracts etc. Various researchers have proved that nutraceuticals are having an important role in reducing the risk of various diseases such as diabetes, cardiovascular diseases, Parkinson's disease etc. with very few or negligible side effects.

Keywords: Nutraceuticals; drugs; dietary supplements; functional foods.

1. INTRODUCTION

"Let food be your medicine and medicine be your food" this great line said by the great scientist,

Hippocrates, is related to the chemicals in food having both nutritional and therapeutic values i.e. Nutraceuticals [1].

*Corresponding author: E-mail: vijaysrampur@gmail.com;

Nutraceuticals are a heterogeneous products category which has a number of synonyms that are used internationally. The term "Nutraceutical" was given by Stephen De Felice who was the founder and chairman of the Foundation for Innovation in Medicine. This term has been part of the industry lexicon for almost a decade [2]. The term "Nutraceutical" is derived from the combination of two words, "nutrient" which means a nourishing food or food component and second "pharmaceutical" which means a medical drug. Nutraceuticals may contain substances that are "natural", intended to treat or prevent a number of diseases, but may not be generally recognized as safe. Hence these are the food products intended for health and medical benefit [3,4]. It has been proved by research that nutraceuticals are useful in providing protection from a number of diseases like diabetes, cancer, cardiac disease, hypertension etc., for example, carotenoids and anti-oxidants found in carrots help in protection against chronic diseases, by preventing free radical damage [5]. Nutraceuticals have become more popular in the modern society due to the belief of reduced chances of adverse effects, being natural, the ease of self-medication and the positive effects on the aging-population.

2. TYPES OF NUTRACEUTICALS

2.1 Dietary Supplements

Dietary supplements (DS) include herbs, minerals, vitamins, or products obtained from plant sources, animal sources such as yeasts, fungus, algae, seafood and many more, for example, energy bars, amino acids, and liquid supplements. They are not consumed in large quantities but have the basic objective to provide nutrition. The United States authorities state that dietary supplements may be regarded as foods, while elsewhere they are classified as drugs or other products [6,7].

2.2 Functional Foods

Japan introduced the concept of functional food in 1980s, to promote health or reduce the risk of diseases. The functional foods include those food items which are advised to be consumed as part of the normal diet, they contain biologically active constituents offering the potential to enhance health or reduce risk of various diseases. Among these foods are those that contain fatty acids, vitamins, specific minerals or dietary fibers, foods with added biologically

active substances such as phytochemicals or other antioxidants and probiotics that have beneficial live cultures [8]. Some examples of functional food products are; milk, cheese and eggs (enriched with omega-3 fatty acids); yogurt enhanced with live active cultures (probiotics); fruit juices and drinks (having antioxidant properties or containing antioxidants); cereals and grains such as wheat, oat, barley (having enriched amounts of dietary fibre); modified fatty acid vegetable oils; soy, canola and hemp (vegetable proteins) and legumes [9,10].

In accordance with the established requirement for the functional food in Japan, the functional food should be consumed:

- a). In its natural form, rather than a prepared dosage form like capsule, tablet, or powder;
- b). Daily, in sufficient quantity; and
- c). In the correct way such that it can regulate a biological process, in order to prevent or cure a disease [11].

2.3 Dietary Supplements and Food Additives

Dietary supplements provide nutrients that may otherwise not be consumed in sufficient quantities. Generally, dietary supplements include vitamins, minerals, fatty acids, fiber or amino acids etc., which according to United State authorities are regarded as foods [12].

Food additives like the dietary supplements, are any substances that are either deliberately added to food to enhance its shelf-life, nutrition, texture, or other quality aspect, or which unintentionally contaminates food (indirect additive) (Fig. 1) [13].

Nutrients are the nutritive constituents present in food that a person consumes for his/her survival and growth. Macronutrients provide the bulk energy required for functioning of metabolic system, whereas micronutrients are helpful in providing the necessary co-factors for metabolism. Plenty of these nutrients are available in the environment [14]. Inorganic chemical compounds like water, minerals, and oxygen should also be considered as nutrients [15].

2.4 Herbals

In ancient times, a large number of herbs were used to prevent and treat many diseases. A plant

containing non-nutritive phytochemicals, provides health benefits if included in diet [16-18]. Nutraceuticals as herbals are big boon to human beings in the aspect of improving their health and to protect them from chronic diseases e.g. willow bark (*Salix Nigra*) helps as anti-inflammatory, analgesic, anti-arthritis, astringent as well as antipyretic [19].

2.4.1 Phytochemicals

Phytochemicals are the plant components having bioactivities which are used to get health benefits, but their use always requires to be backed up with some scientific rationale for being present in food as potential nutraceutical. Phytochemical have following health benefits:

- (1) These are used as substrates for biochemical reactions.
- (2) These are used as cofactors of enzymatic reactions.
- (3) These act as ligands which agonize or antagonize cell surface or intracellular receptors.
- (4) These are used as scavengers of various reactive or toxic chemicals.
- (5) These are used as compounds to increase the absorption and or improve stability of many essential nutrients.

- (6) These work as selective growth factors for gut friendly bacteria.
- (7) Fermented phytochemicals are beneficial for non-pathogenic bacteria found in gastrointestinal tract.
- (8) These are selective inhibitors of deleterious intestinal bacteria. Phytochemicals like terpenoids, phenolics, alkaloids and fiber, are extensively examined and used for their ability to provide health benefits [20].

2.5 Probiotic/ Prebiotics

Probiotic bacteria are “living microorganisms that taken in tolerable quantity, provide a variety of benefits to host health”. These may interact with commensal bacteria to have a direct impact on the host [21]. Metchnikoff was the first person who successfully revamped the toxic flora of the large intestine into a host-sympathetic colony of *Bacillus bulgaricus*, found by Hord [22;23]. Hence probiotics are gut friendly bacteria which aid in digestion and absorption of some nutrients. They act by eliminating the disease causing pathogens, like yeasts, other bacteria and viruses which mutually develop advantageous symbiosis within gastrointestinal tract (Table 1) [24]. The Japanese were the first to recognize the value of non-digestible oligosaccharides, and added these in feed of piglets to relieve and




Fig. 1. Diagrammatic representation of nutraceuticals

prevent diarrhoea. It was observed that fructo-oligosaccharides and galacto-oligosaccharides cause an increase in intestinal bifido-bacteria which stimulated their growth in the human gut. Hence a prebiotic is “a selectively fermented ingredient that causes changes in the activity of the gastrointestinal microbiota that is beneficial for human health” [25]. Some people use lactobacillus for irritable bowel syndrome, Crohn's disease, inflammation of the colon, necrotizing enterocolitis in babies born

prematurely, infection with *Helicobacter pylori*, urinary tract infections, vaginal yeast infections, in prevention of common cold in adults and to prevent respiratory infections in children attending daycare centers. It is also being tested to prevent serious infections in people on ventilators [26-31].

A number of commercial/commercially prepared nutraceuticals are available in market as mentioned in Table 2.

Table 1. Different species of microbes used as probiotic

Genus	Species
<i>Lactobacillus</i>	Acidophilus, Delbrueckia, Brevis, Fermentum, Gasserijohnsonii, Paracasei, Plantarum, Reuteri, Rhamnosus, Salivarius
<i>Bifidobacterium</i>	Adolescentis, Animalisb, Breve, Bifidum, Infantis, Longum
<i>Streptococcus</i>	Thermophilus, Salivarius
<i>Saccharomyces</i>	Cerevisiae
<i>Escherichia</i>	Coli
<i>Enterococcus</i>	Faecium
<i>Bacillus</i>	Coagulans, Clausii

Table 2. List of marketed nutraceuticals

Products	Category	Contents	Manufacturer
Alamin SE	Protein supplement	L-Arginine & other Protein	Albert David Ltd., India
Albumen Care	Protein supplement	L-arginine, multivitamins & minerals	B.V. Bio-Corp Pvt. Ltd., India
Arginine	Protein Supplement	L-arginine, proantho- cyanidis	ManikindPharmaPvt. Ltd, India
Appetite Intercept™	Appetite Suppressant	Caffeine, tyrosine and Phenylalanine	Natrol, Chatsworth, CA, USA
Betafactor™	Immune supplement	Beta glycan	AmeridanInternational Inc. USA
Brainspeed Memory®	Brain Health supplement	Vitamin and minerals	Natrol, Chatsworth, CA, USA
Biovinca™	Neurotonic	Vinpocetine	Cyvex nutrition, Irvine, USA
Coral calcium	Calcium supplement	Calcium and trace minerals	Nature's answer, Hauppauge, NY, USA
Calcirol D-3®	Calcium supplement	Calcium and vitamins	Cadilla healthcare, India
Chaser™	Hangover Supplement	Activated calcium carbonate, and vegetable carbon	Living essentials, Walled lake MI, USA
GRD®	Nutritional Supplement	Proteins, vitamins, minerals and carbohydrates	ZydusCadila Ltd. Ahmedabad, India
Weight smart™	Nutritional supplement	Vitamins and trace elements	Bayer corporation, Morristown, NL, USA
Yakult	Probiotic	Skimmed milk, <i>Lactobacilli</i>	Danoneindia Ltd. India

Products	Category	Contents	Manufacturer
	dairy product	<i>casai Shirota</i>	
ImmunAge	Fermented papaya Preparation	Papaya, yeast, dextrose	Osato laboratory Inc. USA
Glowelle®	Beauty drink	Antioxidants, vitamins and fruit extracts	Nestle, India
HiOwNa	Nutriional supplement	Protein, multivitamins, minerals and antioxidant	Himalaya herbal ltd, India
PediaSure®	Nutritional supplement	Protein, multivitamins and antioxidant	Abbott India ltd, India
Revital®	Health supplement	Ginseng, vitamin and minerals	Ranbaxy, India
Proteinex®	Protein Supplement	Predigested proteins, vitamins, minerals and carbohydrates	Pfizer Ltd., Mumbai, India
Rox® Glucon-D Glucose-D	Energy drink	Taurine, caffeine and glucuronolactone Glucose	Rox America, Spartanburg, SA, USA Dabur
Omega woman	Immune supplement	Antioxidants, vitamins and phytochemicals (eg. Lycopene, and resveratrol)	Wassen, Surrey, U.K
Mushroom optimizer™	Immune Supplement	Mushrooms polysaccharides and Folic acid	Jarrow formulas, Los Angeles, CA, USA
Proplus®	Nutritional Supplement	Soy proteins	Campbell soup company, Camden, NJ, USA
Snapple-aday™	Meal replacement Beverage	Vitamins and minerals	Snapple beverage group, White Plains, NY, USA
WelLife®	Amino acid Supplement	Granulated-L-glutamine	Daesang America Inc., Hackensack, NJ, USA
Oliveinol™	Dietary Supplement	Natural antioxidant, Hydroxytyrosol	CreAgri, Hayward, CA, USA
Threptin®	Diskettes Protein supplements	Proteins and vitamin B	Raptakos, Brett & Co. Ltd., Mumbai, India
Red bull®	Energy drink	Taurine, Caffeine, Glucuronolactone, b- group vitamins	Austrian red bull GmbH

3. PHARMACOLOGICAL NUTRACEUTICALS

Better life quality is achieved with food items filled with nutrient values and remarkably potent in preventing diseases which may be cancer, diabetes, heart diseases, hypertension etc (Table 3 & Fig. 2). Such products may be dietary supplements, food additives, phytochemicals, prebiotics, probiotics, genetically modified food, processed food and plant products.

USE OF 3.1 In Cardiovascular Disease

It is not easy to set up a clear impact of nutrition/physical exercise on major cardiovascular diseases because history of cardiovascular diseases is too long [32]. The effect of calcium on hypertension and pre-eclampsia (a condition in pregnancy characterized by high blood pressure, sometimes with fluid retention and proteinuria) is unpredictable as well as ambiguous. It is

supposed that high levels of intracellular calcium may increase vascular smooth muscle tone, peripheral vascular resistance, and responsiveness to blood pressure. Treatment with vitamin C and selenium need further study to observe its effect on mortality [33-34].

It has been observed that some nutraceuticals may be useful to prevent the risk of thrombosis in women with thrombophilic gene mutations like vitamin E which causes inhibition of platelet aggregation by a protein kinase C-dependent pathway [35-36]. Nutraceuticals like vitamins, minerals, omega-3 poly-unsaturated fatty acids (n-3 PUFAs), dietary fibers and antioxidants, in addition to physical exercise, are recommended to prevent and treat cardiovascular diseases. Researchers have proved that polyphenols found in grapes and in wine are helpful in reducing arterial disease by altering cellular metabolism and signalling [37].

Onion, black grapes, cherries, cruciferous vegetables, grapefruits, red wine, apples and berries are good sources of flavonoids [38] and also available as flavones and flavonols which are beneficial for the treatment of cardiovascular diseases [39-41]. Flavonoids block the angiotensin-converting enzyme, a key moiety that causes a raise in blood pressure [42]. Antioxidant activity of ascorbic acid, alpha-tocopherol, and beta-carotene as has been studied and reviewed [43].

3.2 In the Management of Diabetes

Various *in-vitro* and *in-vivo* studies (animal) have proved that plant polyphenols including phenolic acids, stilbenes, lignans and flavonoids are effective nutraceuticals in diabetes and its prevention, although human clinical trials are required to check the efficacy of poly-phenol compounds in treatment of diabetes [44].

Vitamin C (ascorbic acid) is a chain-breaking antioxidant which prevents the propagation of chain reactions that may lead to a reduction in protein glycation. It has been reported that ascorbic acid helps in reducing diabetes-induced sorbitol in animals [16]. Ascorbic acid supplementation (800 mg/day) partially replenishes ascorbic acid levels in patients with type 2 diabetes but endothelial dysfunction or insulin resistance is not effected at all [17].

Bunyapraphatsara et al. [45] observed a combined effect of *Aloe vera* juice with

glibenclamide in diabetic patients and concluded that glibenclamide alone did not show any effect while *Aloe vera* juice showed significant reduction in fasting blood glucose level and triglycerides within two weeks and four weeks respectively. *Aloe vera*, however, did not show any effect on level of cholesterol, but was found to be effective in the treatment of diabetes [45]. Acacia is non-starch polysaccharides which is not digested in the intestine, but generate short chain fatty acids in large bowel that produce extensive biological effects. Philips et al. [46] conducted study over the extract of Acacia Arabica and confirmed the anti-diabetic effect of acacia resulted by increasing the insulin release. Hou et al. [47] demonstrated significant hypoglycemic effects of Acacia Arabica powder in healthy rabbits, whereas no significant decrease in blood sugar was observed in the alloxan-induced diabetic rabbits was observed. Wadood et al. [48] concluded that Acacia arabica initiates the release of insulin from pancreatic beta cells of normal rabbits. Antidiabetic activity of glycyrrhiza in non-insulin dependent diabetic model was observed by Takii [49].

High intake of isoflavone (20–100 mg/day) is helpful in lowering of rate of mortality in diabetes of type 2, osteoporosis, cardiac disease and certain cancers [50]. Docosahexaenoic acid is vital for neuro-visual development which helps in modulating insulin resistance and nurture the advocacy for essential fatty acids in pregnancy in women with gestational diabetes mellitus [51].

Omega-3 fatty acids helps in reducing blood glucose tolerance in patients predisposed to diabetes. Insulin is needed to synthesize long chain n-3 fatty acids; thus heart may be victim to their depletion in case of diabetes. Hence it is concluded that ethyl esters of n-3 fatty acids may be advantageous in diabetic patients [52]. Lipoic acid is a well-known antioxidant, and in Germany being used to cure diabetic neuropathy. Lipoic acid; as a long-term dietary supplement is aimed at the prophylactic protection of diabetics from complications [53]. α -Lipoic acid enhances insulin sensitivity by approximately 18–20% in patients suffering from type 2 diabetes [54]. Clinical trial studies on α -lipoic acid reported advantageous in the treatment of diabetic neuropathy [55].

Dietary fibers obtained from psyllium have been used to reduce weight and lipid levels in hyperlipidemia as dietary supplement [56]. Intake of Chromium supplements may be helpful to




Fig. 2. Pharmacological uses of nutraceuticals

enhance sensitivity to insulin and boost glucose tolerance in type II diabetic patient [48]. Magnesium-rich diet intake may reduce risk of diabetes by improving in insulin sensitivity [57]. Diabetes management is supported by one of the nutraceuticals Biotin which increase insulin production and stimulates liver glucokinase activity, thus improves the uptake of glucose in muscle cells [58-60]. Pharmacological effect of epinephrine can be inhibited by *Azadirachta Indica* which results in enhanced utilization of peripheral glucose [61,62] and reduce hypoglycaemic activity without change in the serum cortisol level [63,64].

Kernels of *Eugenia jambolana* (*Syzygiumcumini*) are useful in diabetes management; their aqueous/alcoholic extract shows hypoglycemic effect [65]. Green Tulsi (*Ocimumsanctum*) leaves extract also reduces blood sugar significantly by cortisol inhibiting potency as proved in both normal and alloxan induced diabetic rats [66,67].

3.3 In Parkinson's Disease

Latif *et al* concluded that diet enriched with vitamin E may decreases the chances of Parkinson's disease [68] while Brower V reported that creatine is helpful in management of Parkinson's disease by decreasing the clinical symptoms [69].

Antioxidant vitamin supplements such as tocopherol, ascorbic acid and beta-carotene are the abundantly occurring nutraceuticals. As per various earlier literature vitamin E supplements

are becoming popular in treatment of Parkinson's disease, whereas epidemiological studies reported that vitamins C and E rich diets are associated with decreased risk of Parkinson's disease [70-71].

3.4 In Alzheimer's Disease

Literatures have been reported that fulvic acid, an active principle of *Shilajit* is highly effective against brain disorders exclusively and in combination with vitamin B complex. [72] Patients suffering from Alzheimer's disease treated with donepezil and vitamin E found effective although future study was suggested to check and compare additive as well as individual effect [73]. Wettstein et al. [74] reported that mild to moderate Alzheimer's dementia could be treated with metrifonate, donepezil, rivastigmine which are second-generation cholinesterase inhibitors.

Hager et al. [75] found stable cognitive function especially in those patients who were administered with 600 mg Alpha-lipoic acid along with acetyl-cholinesterase inhibitors, in comparison to those patients who only received therapy of standard acetyl-cholinesterase inhibitors since last 337 days. Huperzine-alpha is a plant alkaloid derived from club moss plant (*Huperziaseerrata*), which is a member or the *Lycopodium* species. Huperzine-alpha is in phase III clinical trial in the USA and is available as a dietary supplement [76]. The meta-analysis of Huperzine A reported here highlights that this treatment has certain

significant improvement for patients with Alzheimer's disease and Vascular Dementia, and longer durations may result in better efficacy for patients with Alzheimer's disease [77].

Literature survey supports that Lipoic acid also helps to improve potential of mitochondrial membrane, memory loss due to ageing and brain ailments as well as in patient suffering from Parkinson's and Alzheimer's disease [78].

Table 3. Detailed review on various nutraceuticals

Types of Nutraceuticals	Sources	Active Constituents	Applications
Dietary fibre	Whole grain foods wheat and corn bran, nuts	Insoluble fibre	Reduce chances of colon or breast cancer (anticancer) [94,95], maintain health of digestive tract [96]
	Oats, barley	Beta-Glucan	Reduce risk of cardiovascular disease, lower down Low Density Lipids and total cholesterol [97,98]
	Beans e.g. Legumes, oats, barley and some fibrous fruits	Soluble fibre	Anticancer (Colon Cancer), Digestive [99,100]
Fatty acids	Salmon and other fish oils	Long chain omega-3 Fatty Acids-DHA/EPA	Reduce risk of CVD, Improve mental, visual functions[101-103]
	Cheese, meat products	Conjugated Linoleic Acid	Improving of body composition, Decreases chances of certain cancers [104-106]
	Fruits	Anthocyanidins	Antioxidant; reduce risk of cancer [87-109]
	Green Tea	Catechins	Antitumor [110]
Phenolics	Citrus	Flavonoids	Antioxidative activity, Prevention of coronary heart disease, hepato-protective, Effective in inflammation and cancer [111]
	Cocoa, Chocolate, Cranberries & cranberry	Tannins	Anti-microbial, Reduce risk of cardiovascular disease [112]
	Corn, soy, wheat, wood oils	Plant Sterols, Stanol ester	Lower blood cholesterol levels by inhibiting cholesterol absorption [113]
	Tomatoes	Lycopene	Antioxidant, protect against prostate cancer [114]
Carotenoids	Corn, various fruits, egg yolk, spinach	Lutin	Antioxidant, Muscle regeneration, anti cancer activity, protect eyes against age related muscular degenerations, cataract [115,116]
	Carrots, various fruits (Guava, papaya, Water melon etc) and vegetables (tomatoes etc).	Beta carotene	Antioxidant, protection of cornea against UV light.
	Soya beans	Saponins	Effective against colon cancer, reduces cholesterol level [117]
Probiotics/ Prebiotics	Curd	Lactobacillus	Antibacterial, acute diarrhea [118]
	Whole grains, onions,	Fructo-	Improve GI health, restore gut flora

Types of Nutraceuticals	Sources	Active Constituents	Applications
	combination of Pro & Prebiotics	oligosaccharides	[119]
	Grains	Tocotrienols and tocopherols	The growth of diverse tumors cell lines was suppressed via initiation of apoptosis and concomitant arrest of cells in the G1 phase of the cell cycle [120]
Phytochemical	Cereal grain, dairy & egg products and plants oil	Phytosterols	Exhibit antioxidant, anti-inflammatory, anti-neoplastic, anti-pyretic & immune- modulating activity, decrease cholesterol [121-123]
	Various plants, whole grain	Phenolic constituents	Antioxidants, Anti-hyperglycemic, and anti hypertensive [124]
	Grapes, berries, cocoa, green tea, acacia spp.	Catechin&gallic acids	Antioxidants, Antiradical property, cyto-protective. [125,126]
	Soybeans	Isoflavonoids	Treating cancers & attenuates bone loss [127,128]

3.5 In Erectile Dysfunctioning

L-arginine in combination with pycnogenol, a product obtained from the pine bark (*Pinuspinaster*), is found safe and effective in mild to moderate erectile dysfunction in Japanese patients [79]. When Patients suffering from moderate to severe ED and dyslipidemia were kept on Niacin rich diet, significant improvement was observed in patients [80].

Kaempferia Parviflora Wall Ex. Baker (KP), which is a Thai plant with name, Kra-Chai-Dum and rhizomes of which are used as a traditional medicine to alleviate male impotency, improve male libido, as energizer, control blood pressure and also reduce stomachache. Study reveals that KP is a potential nutraceutical compound effective in male erectile dysfunction caused due to ageing [81].

3.6 Osteoarthritis

Chondroitin sulfate and Glucosamine both are commonly used to alleviate/reduce the symptoms of osteoarthritis. Chondroitin sulfate and Glucosamine both act as nutraceuticals and possible mechanism of their anti- inflammatory activity may be due to synthesis of NO and PGE2 [82]. Capsaicin reduces pain and stiffness and increases joint functioning by acting as agonist for transient receptor potential vanilloid 1 (pain receptor) [82]. Boswelliaserrata relieved joint pain, reduced joint swelling, and stiffness by

inhibiting TNF- α -induced MMP-3 expression and protected against IL-1 β -induced chondrocyte death [83-85]. Capsaicin reduces pain and stiffness to increased joint function by agonising transient receptor potential vanilloid 1 (pain receptor) while prolonged exposure of capsaicin leads to desensitization of this pain pathway [86-87].

Cat's claw reduces osteoarthritis associated pain by Inhibiting lipo-polysaccharide induced PGE2 production and activation of TNF- α . Avocado/soybean unsaponifiables reduced pain in osteoarthritis patients and reduces NSAID consumption by suppressing TNF- α , IL-1 β , COX-2, and iNOS in LPS-activated chondrocytes [88,89]. Collagen hydrolysates alleviate osteoarthritis related pain by stimulating regeneration of type II collagen and by increasing biosynthesis of proteoglycans [90].

3.7 Nutraceuticals Tailoring Genes-Nutrigenomics

Nutrigenomics is an emerging field to know interactions between food and genes, due to manipulated diet [91]. Nutrigenomics is the study of the use of functional genomic tools to probe a biological system followed by a nutritional stimulus to understand how nutritional molecules may affect metabolic pathways and homeostatic control [92]. Study of effect of nutrigenomics also becomes necessary to find out the effect of foods on factors that may interact with particular genes

to increase risk of diseases like diabetes mellitus, obesity, cardiovascular diseases etc. [93]. Hence it is a vast field that describes the impact of food on genes of human being and requires that a lot of studies be done.

4. CONCLUSION

In the present scenario nutraceuticals have become more popular in modern society and became important due to increasing applications of Nutraceuticals which serves as a part of growing pharmaceutical industry. Modern society now is being aware of the food products that are beneficial to them in the aspects of health and nutrition owing to very few or no side effects. Nutraceuticals thus appear to be the way forward to prevent, control and possibly cure chronic diseases in the most natural, safe and easily affordable manner.

CONSENT

It is not applicable.

ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Bagchi D. Nutraceuticals and functional foods regulations in United States and around the world. *Toxicol.* 2006;221:1-3.
2. Granato H. Regulatory concerns cloud functional food. *Nutraceutical Markets*; 2009.
Available:<http://www.naturalproductsinsider.com/articles/2000/11/regulatory-concerns-cloud-functional-food-nutraceutical.aspx>
3. Ross S. Functional foods: The food and drug administration perspective. *Am J Clin Nutr.* 2000;71:1735-38.
4. Palthur MP, Palthur SSS, Chitta SK. Nutraceuticals: Concept and regulatory scenario. *Int J Pharm Pharm Sci.* 2010;2: 14-20.
5. Cindy Oliveri S. Nutraceuticals, phytochemicals, and antioxidants- What are they all about. *OSU Extension Fact Sheet.* 2003;5051-98.
6. Dietary Supplements What You Need to Know, USFDA; 2006.
Available:<http://www.fda.gov/downloads/Food/DietarySupplements/UCM240978.pdf> (Last cited on Jan 20, 2016)
7. Dietary supplement health and education act 1994, public law 103-417, 103rd congress.
Available:http://ods.od.nih.gov/pdf/About/DS_HEA_Wording.aspx.pdf (Last cited on Jan 20, 2016).
8. Available:<http://www.eufic.org/article/en/expid/basics-functional-foods/> (Last cited on Jan 20, 2016)
9. Hathcock J. Dietary supplements: How they are used and regulated. *J Nut.* 2001;131:1114-7.
10. Dureja H, Kaushik D, Kumar V. Developments in nutraceuticals. *Ind J Pharmacol.* 2003;35:363-72.
11. Hardy G. Nutraceuticals and functional foods: Introduction and meaning. *Nutrition.* 2000;16(7-8):688-9.
12. Grace, Emily. How to choose the best supplement. *HealthBeacon.* (Accessed on 3 March 2016)
13. Camire ME, Kantor MA. Dietary supplements: Nutritional and legal considerations. *Food Technol.* 1999;53(7): 87-96.
14. Whitney Elano, Sharo R. Understanding nutrition. 10th ed. Thomson-Wadsworth; 2005.
15. Frances S, Ellie W. Nutrition: concepts and controversies. 1st ed. Cengage Learning; 2007.
16. Hamid AA, Luan YS. Functional properties of dietary fiber prepared from defatted rice bran. *Food Chem.* 2000;68:15-9.
17. Mermel V. Old paths new directions: Use of functional foods in the treatment of obesity. *Trends Food Sci Tech.* 2004;15: 532-40.
18. Dulloo AG, Duret C, Rohrer D. Efficacy of a green tea extract rich in catechin polyphenols and caffeine in increasing 24-h energy expenditure and fat oxidation in humans. *Am J Clin Nutr.* 1999;70:1040-5.
19. Ehrlich SD. (Willow bark), private practice specializing in complementary and alternative medicine Phoenix AZ Review. *VeriMed Healthcare Network*; 2008.
20. Cora JD, German JB. Phytochemicals: Nutraceuticals and human health. *J Sci Food Agric.* 2000;80:1744-56.
21. FAO/WHO. Joint FAO/WHO Expert Consultation on evaluation of health and

- nutritional properties of probiotics in food including powder milk with live lactic acid bacteria Cordoba, Argentina; 2001.
22. Hord NG. Eukaryotic microbiotic crosstalk: potential mechanisms for health benefits of prebiotics and probiotics. *Annu Rev Nutr.* 2008;28:215–31.
23. Michail S, Sylvester F, Fuchs G, Issenma R. Clinical efficacy of probiotics: Review of the evidence with focus on children, clinical practice guideline. *J Pediatr Gastroenterol Nutr.* 2006;43(4):550-7.
24. Holzapfel WH, Haberer P, Geisen R, Bjorkroth J, Schillinger U. Taxonomy and important features of probiotic microorganisms in food and nutrition. *Am J Clin Nutr.* 2001;73:365S-73S.
25. Gibson GR. Dietary prebiotic: Current status and new definition. *IFIS Func Foods Bull.* 2011;7:1–19.
26. Nikfar S, Rahimi R, Rahimi F, et al. Efficacy of probiotics in irritable bowel syndrome: A meta-analysis of randomized, controlled trials. *Dis Colon Rectum.* 2008;51:1775–80.
27. McFarland LV, Dublin S. Meta-analysis of probiotics for the treatment of irritable bowel syndrome. *World J Gastroenterol.* 2008;14:2650-61.
28. AlFaleh K, Anabrees J. Probiotics for prevention of necrotizing enterocolitis in preterm infants. *Cochrane Database Syst Rev.* 2014;10:4.
29. Hamilton-Miller JMT. The role of probiotics in the treatment and prevention of *Helicobacter pylori* infection. *Int J Antimicrob Agent.* 2003;22(4):360–66.
30. Stapleton AE, Au-Yeung M, Hooton TM. Randomized, placebo-controlled phase 2 trial of a *Lactobacillus crispatus* probiotic given intravaginally for prevention of recurrent urinary tract infection. *Clin Infect Dis.* 2011;52(10):1212-17.
31. Strus M, Kucharska A, Kukla G, Brzychczy-Włoch M. The *in vitro* activity of vaginal *Lactobacillus* with probiotic properties against *Candida*. *Inf Dis Obstet Gynecol.* 2005;13(2):69-75.
32. Ignarro LJ, Balestrieri ML, Napoli C. Nutrition, physical activity, and cardiovascular disease: An update. *Cardio Res.* 2007;73:326-40.
33. Trumbo PR, Ellwood KC. Supplemental calcium and risk reduction of hypertension. *Nutr Rev.* 2007;65:78-87.
34. Hatton DC, Yue Q, McCarron DA. Mechanisms of calcium's effects on blood pressure. *Semin Nephrol.* 1995;15(6):593-602.
35. Ravi SMT. Nutrigenetics and nutraceuticals: The next wave riding on personalized medicine. *Trans Res.* 2007; 149:55-61.
36. Freedman JE, Farhat JH, Loscalzo J, et al. Alpha-tocopherol inhibits aggregation of human platelets by a protein kinase C-dependent mechanism. *Circulation.* 1996;94:2434–40.
37. German JB, Walzem RL. The health benefits of wine. *Ann Rev Nutr.* 2000;20: 561-93.
38. Hollman PCH, Hertog MGL, Katan MB. Analysis and health effects of flavonoids. *Food Chem.* 1996;57:43-6.
39. Yao LH, Jiang YM, Shi J, Tomás-Barberán FA, Datta N, Singanusong R, Chen SS. Flavonoids in food and their health benefits. *Plant Foods Hum Nutr.* 2004; 59(3):113-22.
40. Cook NC, Samman S. Flavonoids - Chemistry, metabolism, cardio-protective effects, and dietary sources. *J Nutri Biochem.* 1996;7:66-76.
41. Hollman PCH, Feskens EJ, Katan MB. Tea flavonols in cardiovascular disease and cancer epidemiology. *Proc Soc Exper Biol Med.* 1999;220:198-202.
42. Rajasekaran A, Sivagnanam G, Xavier R. Nutraceuticals as therapeutic agents: A Review. *Res J Pharm Tech.* 2008;1(4): 328-40.
43. Niki E, Noguchi N, Tsuchihashi H, Gotoh N. Interaction among vitamin C, vitamin E, and beta-carotene. *Am J Clin Nutr.* 1995;62(6Suppl):1322S-6S.
44. Zahra B, Parvin M, Fereidoun A. Dietary polyphenols as potential nutraceuticals in management of diabetes: A review. *J Diabetes Metab Disord.* 2013;12(1):43.
45. Bunyapraphatsara N, Yongchaiyudha S, Rungpitarangsi V, Chokeychaijaroenporn O. Antidiabetic activity of Aloe vera L juice II Clinical trial in diabetes mellitus patients in combination with glibenclamide. *Phytomed.* 1996;3(3):245-8.
46. Philips AO, Philips GO. Biofunctional behavior and health benefits of a specific Gum Arabic. *Food Hydrocoll.* 2011;25(2): 165-9.
47. Hou CC, Lin SJ, Cheng JT, Hsu FL. Antidiabetic dimericguianolides and a lignan glycoside from *Lactuca indica*. *J Nat Prod.* 2003;66(5):625-9.

48. Wadood A, Wadood N, Shah SA. Effects of *Acacia arabica* and *Caralluma edulis* on blood glucose levels of normal and alloxan diabetic rabbits. *J Pak Med Assoc.* 1989;39(8):208-12.
49. Takii H, Kometani T, Nishimura T, Nakae T, Okada S, Fushikii T. Antidiabetic effect of glycyrrhizin in genetic diabetes *Licorice* in diabetes. *BiolPharma Bull.* 2000;24: 484-7.
50. Brouns F. Soya isoflavones: A new and promising ingredient for the health foods sector. *Food Res Int.* 2002;35:187-93.
51. Thomas B, Ghebremeskel K, Lowy C, Crawford M, Shore RN. Nutrient intake of women with and without gestational diabetes with a specific focus on fatty acids. *Nutrition.* 2006;22:230-6.
52. Sirtori CR, Galli C. Fatty acids and the Omega 3. *Biomed Pharmacother.* 2002;56: 397-406.
53. Coleman MD, Eason RC, Bailey CJ. The therapeutic use of lipoic acid in diabetes: A current perspective. *Envir Toxi Pharmacol.* 2001;10:167-72.
54. Singh U, Jialal L. Alpha-lipoic acid supplementation and diabetes. *Nutr Rev.* 2008;66:646–57.
55. Ziegler D, Reljanovic M, Mehnert H, Gries FA. Alpha-lipoic acid in the treatment of diabetic polyneuropathy in Germany: Current evidence from clinical trials. *Exp Clin Endo Dia.* 1999;107:421–30.
56. Baljit S. Psyllium as therapeutic and drug delivery agent. *Int J Pharm.* 2007;334:1-14.
57. Lau FC, Bagchi M, Sen CK, Bagchi D. Nutrigenomic basis of beneficial effects of chromium (III) on obesity and diabetes. *Mol Cell Biochem.* 2008;3:171–10.
58. McCarty MF. Nutraceutical resources for diabetes prevention— An update. *Med Hypo.* 2005;64:151–8.
59. Dakshinamurti K, Cheah-Tan C. Biotin-mediated synthesis of hepatic glucokinase in the rat. *Arch Biochem Biophys.* 1968;127:17-21.
60. Furukawa Y. Enhancement of glucose-induced insulin secretion and modification of glucose metabolism by biotin. *Nippon Rinsho.* 1999;57:2261-9.
61. Wang ZQ, Zhang XH, Cefalu WT. Chromium picolinate and biotin enhance glycogen synthesis and glycogen synthase gene expression in human skeletal muscle culture. *Diab Res Clin Pract.* 2000;50:395.
62. Chattopadhyay RR, Chattopadhyay RN, Nandy AK, Poddar G, Maitra SK. The effect of fresh leaves of *Azadirachta indica* on glucose uptake and glycogen content in the isolated rat hemi diaphragm. *Bull Cal Sch Trop Med.* 1987;35:8–12.
63. Chattopadhyay RR. Possible mechanism of antihyperglycemic effect of *Azadirachta indica* leaf extract part IV. *Gen Pharmacol.* 1996;27:431–4.
64. Chattopadhyay RR. A comparative evaluation of some blood sugar lowering agents of plant origin. *J Ethnopharmacol.* 1999;67:367–72.
65. Gholap S, Kar A. Hypoglycaemic effects of some plant extracts are possibly mediated through inhibition in corticosteroid concentration. *Pharmazie.* 2004;59:876–8.
66. Sharma SB, Nasir A, Prabhu K, M Murthy PS, Dev G. Hypoglycaemic and hypolipidemic effect of ethanolic extract of seeds of *Eugenia jambolana* in alloxan-induced diabetic rabbits. *J Ethnopharmacol.* 2003;85:201–6.
67. Rai V, Iyer U, Mani UV. Effect of Tulsi (*Ocimum sanctum*) leaf powder supplementation on blood sugar levels serum lipids and tissue lipids in diabetic rats. *Plant Foods Hum Nutri.* 1997;50:9–16.
68. Latif S, Anwar F, Ashraf M, Gilani AH. *Moringa oleifera*: A food plant with multiple medicinal uses. *Phytother Res.* 2007;21: 17-25.
69. Brower VA. Nutraceutical a day may keep the doctor away. *EMBO Reports.* 2005;8: 708-11.
70. Maxwell SRJ. Antioxidant vitamin supplements: Update of their potential benefits and possible risks. *Drug Saf.* 1999;21:253–66.
71. Anderson C, Checkoway H, Franklin GM, et al. Dietary factors in Parkinson's disease: The role of food groups and specific foods. *Mov Disord.* 1999;14:21–7.
72. Carlos C, Gonzalo F, Patricio F, Fernando C, Ricardo M. Can nutraceuticals prevent alzheimer's disease? Potential therapeutic role of a formulation containing shilajit and complex B vitamins. *Arch Med.* 2012;43(8): 699–704.
73. Klatte ET, Scharre DW, Nagaraja HN, Davis RA, Beversdorf DQ. Combination therapy of donepezil and vitamin E in Alzheimer disease *Alzheimer. Dis Asso Disord.* 2003;17:113-6.

74. Wettstein A, Stadtärztlcher DZ. Cholinesterase inhibitors and Ginkgo extracts-- Are they comparable in the treatment of dementia? Comparison of published placebo-controlled efficacy studies of at least six months' duration. *Phytomedicine*. 2000;6(6):393-401.
75. Hager K, Marahrens A, Kenkies M, Riederer P, Munch G. Alphaipoic acid as a new treatment option for Alzheimer type dementia. *Arch Gerontol Geriatr*. 2001;32: 275-82.
76. Grunwald J, Raveh L, Doctor BP. Huperzine A as a pretreatment candidate drug against nerve agent toxicity. *Life Sci*. 1994;54:991-7.
77. Shu-huai X, Chun-Xiao Z, Rui Z, Li An. Huperzine A in the treatment of alzheimer's disease and vascular dementia: A Meta-Analysis. *Evid Based Complement Alternat Med*. 2014;10. Article ID 363985.
Available:<http://dxdoiorg/101155/2014/363985>
78. Kramer K, Elkins MHR. Alpha lipoic acid: Nature's supreme antioxidant. In: Elkins MHR. USA: Wood land Publishing; 1998.
79. Aoki H, Nagao J, Ueda T, Strong JM, Schonlau F, Yu-Jing S, Lu Y, Horie S. Clinical assessment of a supplement of Pycnogenol and L-arginine in Japanese patients with mild to moderate erectile dysfunction. *Phytother Res*. 2012;26(2): 204-7.
80. Ng CF, Lee CP, Ho AL, Lee VW. Effect of niacin on erectile function in men suffering erectile dysfunction and dyslipidemia. *J Sex Med*. 2011;8(10):2883-93.
81. Yenjai CK, Prasanphen S, Daodee V, Wongpanich, Kittakoop P. Bioactive flavonoids from *Kaempferia parviflora*. *Fitoterapia*. 2004;75(1):89-92.
82. Kalioraa AC, Dedoussisa GVZ, Schmidt H. Dietary antioxidants in preventing atherogenesis. *Atherosclerosis*. 2006; 187(1):1-17.
83. Krishnaraju AV, Sundararaju D, Vamsikrishna U, Suryachandra R, Machiraju G, Sengupta, K, Trimurtulu G. Safety and toxicological evaluation of Aflapin: A novel *Boswellia*-derived anti-inflammatory product. *Toxicol Mech Method*. 2010;20:556-63.
84. Kimmatkar N, Thawani V, Hingorani L, Khiyani R. Efficacy and tolerability of *Boswelliaserrata* extract in treatment of osteoarthritis of knee— A randomized double blind placebo controlled trial. *Phytomedicine*. 2003;10:3-7.
85. Sengupta K, Kolla JN, Krishnaraju AV, Yalamanchil N, Rao CV, Golakoti T, Raychaudhuri S, Raychaudhuri SP. Cellular and molecular mechanisms of anti-inflammatory effect of Aflapin: A novel *Boswellia serrata* extract. *Mol. Cell Biochem*. 2011;354:189-97.
86. Kosuwon W, Sirichatiwapee W, Wisanuyotin T, Jeeravipoolvarn P, Laupattarakasem W. Efficacy of symptomatic control of knee osteoarthritis with 0.0125% of capsaicin versus placebo. *J. Med. Assoc*. 2010;93:1188-95.
87. Remadevi R, Szallisi A. Adlea (ALGRX-4975), an injectable capsaicin (TRPV1 receptor agonist) formulation for longlasting pain relief. *Drugs*. 2008;11: 120-32.
88. Christensen R, Bartels EM, Astrup A, Bliddal H. Symptomatic efficacy of avocado-soybean unsaponifiables (ASU) in osteoarthritis (OA) patients: A meta-analysis of randomized controlled trials. *Osteoarthr. Cartilage*. 2008;16:399-408.
89. Au RY, Al-Talib TK, Au AY, Phan PV, Frondoza CG. Avocado soybean unsaponifiables (ASU) suppress TNF-alpha, IL-1beta, COX-2, iNOS gene expression, and prostaglandin E2 and nitric oxide production in articular chondrocytes and monocyte/macrophages. *Osteoarthr. Cartilage*. 2007;15:1249-55.
90. Bello AE, Oesser S. Collagen hydrolysate for the treatment of osteoarthritis and other joint disorders: A review of the literature. *Curr Med Res Opin*. 2006;22:2221-32.
91. Neeha VS, Kint P. Nutrigenomics research: A review. *J Food Sci Technol*. 2013;50(3):415-28.
92. Mutch DM, Wahli W, Williamson G. Nutrigenomics and nutrigenetics: The emerging faces of nutrition. *FASEB Journal*. 2005;1602-16.
93. Nutrigenomics focuses on the effect of nutrients on the genome, proteome, and metabolome. *Ordovas J, Mooser M. Current Opinion in Lipidology*. 2004;15: 101-8.
94. Adom KK, Liu RH. Antioxidant activity of grains. *J Agric Food Chem*. 2002;50(21): 6182-7.
95. Ferguson LR, Harris P. Protection against cancer by wheat bran: Role of dietary fibre

- and phytochemicals. *Euro J Can Pre*. 1999;8(1):17-25.
96. Engberg RM, Hedemann MS, Steenfeldt S, Jensen BB. Influence of whole wheat and xylanase on broiler performance and microbial composition and activity in the digestive tract. *Poul Sci*. 2004;83(6):925-38.
97. Keogh GF, Cooper GJS, Mulvey TB, McArdle BH, Coles GD, Monro JA, Poppitt SD. Randomized controlled crossover study of the effect of a highly β -glucan-enriched barley on cardiovascular disease risk factors in mildly hypercholesterolemic men. *Am J Clin Nutr*. 2003;78(4):711-8.
98. Theuvsen E, Mensink RP. Water-soluble dietary fibers and cardiovascular disease. *Physiol Behav*. 2008;94(2):285-92.
99. McIntyre A, Gibson PR, Young GP. Butyrate production from dietary fibre and protection against large bowel cancer in a rat model. *Gut*. 1993;34:386-91.
100. Malkki Y, Virtanen E. Gastrointestinal effect of oat bran and oat gum: A review. *LWT- Food Sci Technol*. 2001;34(6):337-47.
101. Popp-Snijders C, Schouten JA, Heine RJ, Vander MJ, Vander VE. A Dietary supplementation of omega-3 polyunsaturated fatty acids improves insulin sensitivity in non-insulin-dependent diabetes. *Diabetes Res*. 1987;4(3):141-7.
102. Connor WE. Importance of n-3 fatty acids in health and disease. *Am J Clin Nutr*. 2000;71(1):171S-5S.
103. Young G, Conquer J. Omega-3 fatty acids and neuropsychiatric disorders. *Reprod Nutrition Dev*. 2005;45(1):1-2,8.
104. Lee KN, Kritchevsky D, Pariza MW. Conjugated linoleic acid and atherosclerosis in rabbits. *Atherosclerosis*. 1994;108(1):19-25.
105. Clement I, Banni S, Angioni E, Carta G, McGinley J, Thompson HJ, Barbano D, Bauman D. Conjugated Linoleic Acid-Enriched butter fat alters mammary gland morphogenesis and reduces cancer risk in rats. *J Nutr*. 1999;129(12):2135-42.
106. Shultz TD, Chewab BP, Seamana WR, Luedeck LO. Effect of conjugated linoleic acid on body composition in mice. *Cancer Lett*. 1992;63(2):125-33.
107. Wang H, Cao G, Prior RL. Oxygen radical absorbing capacity of anthocyanins. *J Agric Food Chem*. 1997;45(2):304-9.
108. Kanga SY, Seeramb NP, Narib MG, Bourquin LD. Tart cherry anthocyanins inhibit tumor development in ApcMin mice and reduce proliferation of human colon cancer cells. *Cancer Lett*. 2003;194(1):13-9.
109. Li-Shu W, Stoner GD. Anthocyanins and their role in cancer prevention. *Cancer Lett*. 2008;269(2):281-90.
110. Liang G, Tang A, Lin X, Li L, Zhang S, Huang Z, Tang H, Li QQ. Green tea catechins augment the antitumor activity of doxorubicin in an *in vivo* mouse model for chemoresistant liver cancer. *Int J Oncol*. 2010;37(1):111-23.
111. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: An overview. *Scientific World J*. 2013;1-16.
112. Lim H, Darah I, Jain K. Antimicrobial activities of tannins extracted from *Rhizophora apiculata* barks. *J Trop Forest Sci*. 2006;18(1):59-65.
113. Katan MB, Grundy SM, Jones P. Efficacy and safety of plant stanols and sterols in the management of blood cholesterol levels. *Mayo Clin Proc*. 2003;78(8):965-78.
114. Takeoka GR, Dao L, Flessa S, Gillespie DM, Jewell WT, Huebner B, Bertow D, Ebeler SE. Processing effects on lycopene content and antioxidant activity of tomatoes. *J Agric Food Chem*. 2001;49(8):3713-17.
115. Naguib YMA. Antioxidant activities of astaxanthin and related carotenoids. *J Agric Food Chem*. 2000;48(4):1150-54.
116. Olmedilla B, Granado F, Blanco I, Vaquero M, Cajigal C. Lutein in patients with cataracts and age-related macular degeneration: A long-term supplementation study. *J Sci Food Agri*. 2001;81(9):904-9.
117. Kim HY, Yu R, Kim JS, Kim YK, Sung MK. Antiproliferative crude soy saponin extract modulates the expression of I κ B α protein kinase C and cyclooxygenase-2 in human colon cancer cells. *Cancer Lett*. 2004;210(1):1-6.
118. Isolauri E, Rautanen T, Juntunen M, Sillanauke P, Koivula T. A human *Lactobacillus* strain (*Lactobacillus Caseisp* strain GG) promotes recovery from acute diarrhea in children. *Pediatric*. 1991;88(1):90-7.
119. Fuller R, Gibson GR. Probiotics and prebiotics: Microflora management for improved gut health. *Clinic MicrobiolInfect*. 1998;9:477-80.

120. Ju J, Picinich SC, Yang Z. Cancer-preventive activities of tocopherols and tocotrienols. *Carcinogenesis*. 2010;31(4): 533–42.
121. Wang T, Hicks KB, Moreau R. Antioxidant activity of phytosteroloryzanol and other phytosterol conjugates. *J Ame Oil Chem Soci*. 2002;79(12):1201-6.
122. Rozner S, Garti N. The activity and absorption relationship of cholesterol and phytosterols. *Colloids and Surfaces A: Physicochem Engg Aspec*. 2006;282-283:435-56.
123. Garcia MD, Saenz MT, Gomez MA. Topical anti-inflammatory activity of phytosterol isolated from *Ezyngium foetidum* on chronic and acute inflammatory model. *Phytother Res*. 1999;13(1):78-80.
124. Ranilla LG, Kwon Y, Apostolidis E, Shetty K. Phenolic compounds antioxidant activity and in vitro inhibitory potential against key enzymes relevant for hyperglycemia and hypertension of commonly used medicinal plants herbs and spices in Latin America. *Bioresour Tech*. 2010;101(12):4676–89.
125. Morel I, Lescoat G, Cogrel P, Sergeant O, Padeloup N. Antioxidant and iron-chelating activities of the flavonoids catechinquercetin and diosmetin on iron-loaded rat hepatocyte cultures. *Biochem Pharmacol*. 1993;45(1):13-9.
126. Kroes BH, Vanden BAJ, Quarlesvan UHC, Labadie RP. Anti-inflammatory activity of gallic acid. *Planta Medica*. 1992;58(6):499-504.
127. Birt DF, Hendrich S, Wang W. Dietary agents in cancer prevention: flavonoids and isoflavonoids. *Pharmacol Thera*. 2001;90(2–3):157–77.
128. Alekel DL, St G, Peterson CT, Hanson KB, Stewart, Toda T. Isoflavone-rich soy protein isolate attenuates bone loss in the lumbar spine of perimenopausal women. *Am J Clin Nut*. 2000;72(3):844-52.

© 2016 Sharma et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://sciencedomain.org/review-history/14624>